

RiskRadar mobiliosios aplikacijos (programos) įdiegimo vadovas

DATA	2019/11/25
AUTORIAI:	Christine KAKALOU (CERTH), Vassilis KOUTKIAS (CERTH)
DOKUMENTO VERSIJA:	1.1



Co-funded by the Health Programme of the European Union Finansuojama iš 3-iosios Europos Sąjungos Sveikatos programos pagal dotacijos susitarimą Nr. 761319

Turinys

1.	Mob	iliosios aplikacijos įdiegimas Android įrenginiuose	3
	1.1	Programos įdiegimas	3
	1.2	Naršymo Android programoje principai	4
	1.3	Android programos rizikos skaičiuoklės pavyzdys	6
2.	Mob	iliosios aplikacijos įdiegimas iOS įrenginiuose	.10
	2.1	Programos įdiegimas	.10
	2.2	Naršymo iOS programoje principai	.12
	2.3	iOS aplikacijos rizikos skaičiuoklės pavyzdys	.14

1. Mobiliosios aplikacijos įdiegimas Android įrenginiuose

1.1 Programos įdiegimas

Norėdami įdiegti RiskRadar demo versiją, pirmiausiai turite įsitikinti, kad Jūsų mobiliajam įrenginiui suteikta teisė įdiegti mobiliąsias programėles iš kitų nei Google Play Store šaltinių:

SETTI	tf(76% ₩ 13:22	14:29	& 😇 🚛 46 🎟	14:29	<table-cell> 🖯 🚛 46 🍘</table-cell>
	Sounds and ultration	< Additional setti	ngs	< Pri	vacy
<1))	Sounds, Vibration, Do not disturb	Date & time	>	DEVICE ADMINISTRATION	
1	Notifications Block, allow, prioritipe	Languages & input	>	Device administrators	ministrators
Ģ	Display Brightness, Home screen	Region	Greece >	Unknown sources	
	Advanced features Games, One-handed mode	Privacy	>	Allow installation of apps fro sources	om unknown
	Device maintenance Battery, Storage, Memory	Enterprise mode	2	Location	>
	Apps Default apps, App permissions	Authorization & revocation	>	Notification access	7 apps can read notifications
ę	Wallpapers and themes Wallpapers, Themes, Icons			Apps with usage acces	s >
	Lock screen and security Lock screen, Fingerprints			Device ID	>

Išmaniojo telefono meniu "Nustatymai (angl. Settings)" priklausomai nuo telefono pasirinkę "Sauga (angl. Security)", "Privatumas (angl. Privacy)" ir kt., aktyvuokite mygtuką "Leisti diegti programas iš nežinomų šaltinių (angl. Allow installation of apps from unknown sources)".

	A # 13-23	±	♥ ./ 49% ii 17:01
< LOCK SCREEN AND SECURITY	Q	INTEGRATE Toolkit	INTEGRATE Toolkit
Samsung Pass Lise biometric authentication to verify your identication to verify your identication to verify your identication of security	stay.	Do you want to install this application? It does not require any special access.	
Find My Mobile		в	ocked by Play Protect
Unknown sources			INTEGRATE Toolkit
Installing from unknown sources may be harmful to your device and		PL ag de	y Protect doesn't recognise this p's developer. Apps from unknown relopers can sometimes be unsafe.
personal data. By tapping OK, you agree that you are solely respons- ible for any damage to your device or loss of data that may result froo using these applications.	n		
CANCEL	ок.	CANCEL INSTALL	



RisRadar programos piktograma (po įdiegimo)

Apsilankykite tinklalapyje: <u>https://integrateja.eu/content/riskradar</u> ir pasirinkite Android aplikaciją.

Gali pasirodyti keli įspėjimai. Norėdami tęsti diegimo procesą, spauskite mygtuką "Gerai (angl. OK)".

Įdiegus RiskRadar programą, jos turinys bus automatiškai pritaikytas kalbai (lietuvių, anglų, kroatų) pagal jūsų Android telefoną. Jei telefone nustatyta kita kalba, tuomet turinys bus rodomas anglų kalba.

1.2 Naršymo Android programoje principai



Pirmasis ekranas paleidžiant RiskRadar programą (RR).

🔝 🔌 🧌 62% 🖥 13:18

RiskRadar

for HIV, hepatitis, STIs and TB

Are you a:

00A...

O Man

O Woman

O Trans woman

O Trans man



Paspaudus mygtuką "Toliau", rodomi RR komponentai.





Turinio informacijos šaltinis, rodomas pasirinkus RR komponentą "Faktų suvestinės".



Pirmasis ekranas, rodomas pasirikus RR komponentą "Rizikos skaičiuoklė".



Pirmasis ekranas, rodomas pasirinkus "Faktų suvestinės".



Hepatitis C is a liver disease caused by infection with the hepatitis C virus (HCV). HCV can cause both acute and chronic hepatitis infection, ranging in severity from a mild illness that lasts only a few weeks to a serious, lifelong illness resulting in cirhosis and liver cancer. The virus is mainly acquired by contact through broken skin with infectious blood. In Europe, the main route of HCV transmission is via injecting drug use as a result of sharing contaminated needles. More rarely, the virus can be transmitted sexually, in healthcare settings due to inadequate infection control practices or perinatally from an infected mother to the baby. Most people with acute HCV infection do not have any symptoms. Those who develop chronic infection are often asymptomatic until decades after infection when symptoms develop secondary to serious liver damage. Around 30% of people with chronic hepatits C suffer from liver damage and a small number of those develop

Faktų suvestinių temos pasirenkamos braukiant į abi puses.



Pranešimas, rodomas pasirinkus RR komponentą "Neaptinkamas = neperduodamas".



Informacija apie tuberkuliozę rodoma, pasirinkus RR komponentą "Tuberkuliozė".



Norėdami surasti artimiausią ištyrimo vietą, pasirinkite RR komponentą "Tyrimų paieška". Priklausomai nuo nustatytos kalbos, būsite nukreipti į atitinkamos šalies žemėlapį.

P		tf .il 48% 🗎	17:04
	re-Exposure Pr	ophylaxis	i
	Does PrEP work?	Is PrEP worth it	t? Ir
Antiretro to treat p way of gi to prever People li viral ther normal li now happ they have are alrea ART. But havir combina lives, and have side Although condition People lin rejection they may disclose they pass PrEP, in c HIV drug one pill, a to take it	virals (ARVs) are to iving ARVs to HIV- iving ARVs to HIV- it them getting HI ving with HIV who- apy (ART) can nor fe span; most dee pen in people who a HIV, start treatm dy seriously ill, or ig HIV requires th tion of drugs for t of for some people e-effects. HIV is now a mai and violence. In r also face jail if ti their infection, wh s on HIV or even o contrast, involves or, more often, tw and only for as lor	the drugs used HIV. PrEP is a negative peop V. take antiretro wexpect a nea this due to AII to do not know tent when they who cannot g at people take he rest of thei the drugs will nageable tigmatised on face isolation many countrie ney fail to tether or not could do so. taking only on to HIV drugs in ng as you need	d o- ar- ODS vet ea ir l e. , , s, e n d

Informacija apie PrEP rodoma, pasirinkus RR komponentą "PrEP".



Detali išyrimo vietos informacija rodoma, paspaudus žymeklį, esantį žemėlapyje.

0 🖬		瀏寒 常川 87% ■ 10:26
About	t INTEGRATE	E JA
Find u	us on Social	Media
Conta	ict us	
Remir	nder	

Mobiliosios aplikacijos nustatymai ir kita informacija.

1.3 Android programos rizikos skaičiuoklės pavyzdys







月页

🔊 🛠 🕾 🗃 58% 🛢 13:30

Risk calculator results

You might have been exposed to hepatitis and other STIs; your partner might also have been exposed to HIV. You should both get a full screening, which includes testing for HIV, hepatitis and STIs. Your partners should start taking PrEP to protect themselves from getting infected with HIV via sex.



000

Wit is recommended that all gay and bisexual men get vaccinated for Hepatitis A & B; consider asking your medical doctor for a Hepatitis A & B vaccination (You will need a test before getting vaccinated).

2. Mobiliosios aplikacijos įdiegimas iOS įrenginiuose

2.1 Programos įdiegimas

Norėdami įdiegti RiskRadar programos demo versiją, apsilankykite šiame tinklalapyje: <u>https://integrateja.eu/content/riskradar</u> ir pasirinkite iOS aplikaciją. Jūsų išmaniojo telefono naršyklėje automatiškai atsidarys "TestFlight" puslapis, kuriame turėsite atlikti du žingsnius:



Apsilankykite tinklalapyje: https://integrateja.eu/content/riskrad ar ir pasirinkite iOS aplikaciją.

Jūsų išmaniojo telefono naršyklėje automatiškai atsidarys "TestFlight" puslapis Pirmame žingsnyje "Get TestFlight" spauskite mygtuką "View on App Store", kad galėtumėte atsisiųsti TestFlight aplikaciją.


Puslapis 11 iš 18

2.2 Naršymo iOS programoje principai



RisRadar programos (RR) piktograma (po įdiegimo).



Pirmasis ekranas, rodomas pasirikus RR komponentą "Rizikos skaičiuoklė".



Pirmasis ekranas paleidžiant RR.



Pirmas rizikos skaičiuoklės klausimas.



Paspaudus mygtuką "Tęsti", rodomi RR komponentai.



Turinio informacijos šaltinis, rodomas pasirinkus RR komponentą "Faktų suvestinės".

Constright all 13:26 100% HIV, Hepatitis and STIs HIV/AIDS Hepatitis A Hepatitis B Hepatitis C Chlamyde

The human immunodeficiency virus (HIV) remains one of the most important communicable diseases in Europe. It is an infection associated with serious disease, persistently high costs of treatment and care, significant number of deaths and shortened life expectancy. HIV is a virus which attacks the immune system and causes a lifelong severe illness with a long incubation period.

Transmission

HIV infection is spread by sexual contact with an infected person, by sharing needles or syringes (primarily for drug injection) with someone who is infected, or, less commonly (and now very rarely in countries where blood is screened for HIV antibodies), through transfusions of infected blood or blood clotting factors. Babies born to HIV-infected women may become infected before or during birth or through breast-feeding. The end-stage of the infection, acquired immunodeficiency syndrome (AIDS), results from the destruction of the immune system. AIDS is

Pirmasis ekranas, rodomas pasirinkus "Faktų suvestinės".

TestFlight atl 13:27 100% HIV, Hepatitis and STIs

Hepatitis B is a liver disease that results from infection with the hepatitis B virus (HBV) and is spread through contact with infected body fluids or blood products. Following acute infection with HBV, some people go on to develop a chronic infection.

Transmission routes and symptoms

Hepatitis B is spread through contact with contaminated body fluids. Most infections occur by coming into contact with infected blood but semen, saliva and cervical secretions can also be infectious. The virus can live on surfaces for at least seven days which means it can be transmitted via objects that have been contaminated with infected body fluids (e.g. used needles). Sexual transmission and injecting drug use are the most common current routes of transmission in Europe. Transmission may also occur in healthcare settings due to the reuse or inadequate sterilisation of medical equipment, especially syringes and needles. Transmission via blood transfusion or through the use of plasma-derived products is now rare in Europe due to

Faktų suvestinių temos pasirenkamos braukiant į abi puses.



PrEP works if you take it.

Most studies have shown convincingly that PrEP is highly effective at preventing HIV infection via sexual transmission, in both men who have sex with men (MSM) and heterosexual women and men.

Some studies have found that PrEP had no effectiveness. The lack of effect seen in these studies was entirely due to very low rates of actual use of PrEP.

A meta-analysis by the World Health Organization published in July 2016, found that, averaged over all studies, PrEP stopped seven out of every ten HIV infections that would otherwise have happened. This meta-analysis by definition includes studies with very low rates of PrEP adherence.

If taken consistently enough, PrEP is almost 100% effective. There are only three convincingly documented cases, among well over 100,000 users, where people acquired HIV despite good adherence to PrEP; in two of these cases the person was unlucky enough to acquire a rare, highly drug-resistant virus.

Informacija apie PrEP rodoma, pasirinkus RR komponentą "PrEP".



Pranešimas, rodomas pasirinkus RR komponentą "Neaptinkamas = neperduodamas".



13:27

C TestFlight

Note: The information contained in these fact sheets is intended for the purpose of general information and should not be used as a substitute for the individual expertise and judgement of healthcare professionals.

Informacija apie tuberkuliozę rodoma, pasirinkus RR komponentą "Tuberkuliozė".



Norėdami surasti artimiausią ištyrimo vietą, pasirinkite RR komponentą "Tyrimų paieška". Priklausomai nuo nustatytos kalbos, būsite nukreipti į atitinkamos šalies žemėlapį.



Detali išyrimo vietos informacija rodoma, paspaudus žymeklį, esantį žemėlapyje.

III vf-GR 🗢	13:46	100% 📖
About INTEG	RATE JA	>
Find us on Sc	cial Media	>
Contact us		>
Test Reminde	r	\bigcirc

The INTEGRATE Toolkit has been developed by the Centre for Research & Technology Hellas (CERTH), https://www.certh.gr/ For potential technical issues, please contact hmw@certh.gr



Mobiliosios aplikacijos nustatymai ir kita informacija.

2.3 iOS programos rizikos skaičiuoklės pavyzdys



uil vf-GR 🗢	13:47 Risk Calculator	100%	uil vf-GR ♥	13:47 Risk Calculator	100%	uil vf-GR 🗢	13:48 Risk Calculator	100% 🗯
	Integrate			Integrate			Integrate	
Do Yes No	you have sex with me Continue	en?	Wh I am H My pa Both r I had a I injec	ay are you using this calculator? AIV+ artner is HIV+ me and my partner are is condomless sex with so t drugs st curious Continue	risk HIV+ omeone	Have pr substa Yes No	you ever injected any escribed drugs or oti ances, or had someou inject you? Continue	y non- ter ne else
∎II vf-GR 🗢	13:48 Risk Calculator	100% 🗪	nil vf-GR ♥	13:48 Risk Calculator	100% 📩	uil vf-gr 중	13:48 Risk Calculator	100% 💻
Are you Yes No I don't	Integrate vaccinated for hepa	atitis B?	Were yo	Integrate bu born outside of	Europe?	Have yo sin Yes No I don't	Integrate ou had a sexual exp ce your last check-	erience up?
	Continue			Continue			Continue	







